



Knowledge grows

YaraVita[®] MOLYTRAC[™]

A formulated product for the treatment of molybdenum deficiency by foliar application

Guaranteed Analysis	
Available Phosphate (P ₂ O ₅)	15%
Molybdenum (Mo)	15.5%

The information provided is accurate to the best of Yara's knowledge and belief. Any recommendations are meant as a guide and must be adapted to suit local conditions.



Benefits

- Formulated for safe application at critical growth stages to satisfy crop requirements
- Widely tank mixable with other crop sprays. Visit www.tankmix.com/yara for details.
- Proven, reliable performance. Tried and tested on a wide range of crops around the world
- High quality, consistent product. Manufactured to ISO 9001 quality assurance standards
- Easy to use liquid formulation. Pours and disperses easily and quickly into the spray tank.
- High nutrient content means lower application rates reducing handling time and waste packaging



Product Recommendations

Typical Crop Recommendations*

- **Alfalfa:** 250 to 500 mls/ha every 2 to 3 cuts. Water rate: 50 to 200 l/ha
- **Beans:** 250 ml/ha when plants are 5 to 15 cm tall. For moderate to severe deficiency, repeat the application once or twice at 10 to 14 day intervals. Water rate: 200 l/ha.
- **Brassicas:** 250 ml/ha at the 4 to 6 leaf stage. For moderate to severe deficiency, repeat the application once or twice at 10 to 14 day intervals. Water rate: 200 l/ha.
- **Carrots:** 250 ml/ha when the crop is 15 cm tall. Water rate: 200 l/ha.
- **Cereals (Barley, Wheat, Oats):** 125 to 250 ml/ha pre-plant soil application (applied with herbicide where tank mixable). Water rate: 30 to 100 l/ha.
- **Cucurbits (field grown):** 250 ml/ha 1 to 2 weeks after germination and at the 4 leaf stage. Water rate: 200 l/ha.
- **Oilseed rape (canola):** For a single application, 250 ml/ha at onset of stem extension. For moderate deficiency, 250 ml/ha at 4 to 6 leaf stage and 250 ml/ha at onset of stem extension. An extra application can be made 10 to 14 days later for severe deficiency. Avoid flowering. Water rate: 30 to 200 l/ha.
- **Soybeans:** 250 to 500 ml/ha when plants are 5 to 15 cm tall. Repeat the application once or twice at 10 to 14 day intervals if necessary. Water rate: 200 l/ha.
- **Squash (field grown):** 250 ml/ha 1 to 2 weeks after germination and at the 4 leaf stage. Water rate: 200 l/ha.
- **Sugar beet:** 250 ml/ha at 4 to 6 leaf stage. Repeat the application once or twice at 10 to 14 day intervals if necessary. Water rate: 200 l/ha.
- **Tomatoes (field grown):** 250 ml/ha when crop is 15 cm tall. Water rate: 500 l/ha.

*The information provided is accurate to the best of Yara's knowledge and belief. Any recommendations are meant as a guide and must be adapted to suit local conditions. Always read the label before use.

1700020

