




Knowledge grows



YaraVita[®] HYDROPHOS[™]

A concentrated liquid phosphorus with magnesium and potassium for use on potatoes, corn and legumes.
Guaranteed Analysis: soluble in water

Available phosphoric acid (P ₂ O ₅)	29% w/w
Soluble potash (K ₂ O)	5% w/w
Magnesium (Mg)	4% w/w

YaraVita HYDROPHOS is a highly efficient source of plant-available foliar phosphorus to provide the energy boosts that complement a soil application strategy.

YaraVita HYDROPHOS

- contains 100% plant active ortho-phosphate to provide an instant energy boost to the crop, as well as additional magnesium to boost photosynthetic activity.
- has a low pH formulation that optimizes nutrient delivery and enhances tank mixability with agrochemicals.
- is a liquid formulation that is easy to handle and is widely tank mixable for coapplication with agrochemicals, removing the need for specific spray operations.

Benefits

- Formulated for safe application at critical growth stages to satisfy crop requirements
- Widely tank mixable with other crop sprays. Visit www.tankmix.com for details
- Proven, reliable performance. Trialed and tested on a wide range of crops around the world
- High quality, consistent product. Manufactured to ISO 9001 quality assurance standards.
- Easy to use flowable formulation. Pours and disperses easily and quickly into the spray tank
- High nutrient content means lower application rates reducing handling time and waste packaging
- Widely used for foliar application to potatoes.



Product Recommendations

Apples, Pears:

4 to 6 applications of 5 l/ha (2 l/ac) from end of flowering. Repeat applications at 10-14 day intervals. Water rate: 500 to 1,000 l/ha (202-404 l/ac).

Beans:

3 to 5 l/ha (1.21-2 l/ac) when crop is 15 cm tall. Repeat if necessary at 10 to 14 day intervals. Water rate: 30 to 200 l/ha (12.14-81 l/ac).

Blueberries:

10 l/ha (4 l/ac) 10 days after petal fall has finished. Water rate: 1,000 l/ha (404 l/ac).

Brassicas:

4 l/ha (1.62 l/ac) soon after transplanting (e.g. 10 to 14 days). Water rate: 200 l/ha (81 l/ac).

Carrots:

2 to 3 applications of 5 l/ha (2 l/ac) applied when the crop is 15 cm tall and at 7 to 14 day intervals. Water rate: 200 l/ha (81 l/ac).

Cereals (Barley, Wheat, Oats):

5 l/ha (2 l/ac) at tillering. Repeat at 10 to 14 day intervals if necessary. Also, apply from ear emergence until the end of flowering (Zadok's G.S. 51-69). Water rate: 200 l/ha (81 l/ac).

Cucurbits (Field Grown):

5 l/ha (2 l/ac) at 4 leaf stage, with second application 10 to 14 days later if necessary. Water rate: 500 l/ha (202 l/ac).

Ginseng:

5 l/ha (2 l/ac) once the new season growth is underway. For moderate to severe deficiency repeat applications at 10 to 14 day intervals. Water rate: 500 l/ha (202 l/ac).

Maize (Corn):

5 l/ha (2 l/ac) at the 4 to 6 leaf stage. Repeat at 10 to 14 day intervals if necessary. Water rate: 200 l/ha (81 l/ac).

Canola:

For a single application, 3 to 5 l/ha (1.21-2 l/ac) at onset of stem extension. For moderate deficiency, 3 to 5 l/ha (1.21 - 2 l/ac) at 4 to 6 leaf stage and at onset of stem extension. An extra application can be made 10 to 14 days later for severe deficiency. Avoid flowering. Water rate: 30 to 200 l/ha (12.14-81 l/ac).

Potatoes:

To increase tuber number, 10 l/ha (4 l/ac) at tuber initiation (when 50% of tip swellings are twice the diameter of the rest of the stolon). To increase tuber size, a minimum of 2 applications of 5 l/ha (2 l/ac) during tuber bulking (when the first formed tubers are greater than 10 mm in diameter). Allow 10 to 14 days between applications. Also, apply during tuber bulking following petiole analysis. Water rate: 200 l/ha (81 l/ac).

Raspberry (Field Grown):

3 x 10 l/ha (4 l/ac) at start of flowering, end of flowering, and fruit development. Water rate: 500 l/ha (202 l/ac).

Squash (Field Grown):

5 l/ha (2 l/ac) at 4 leaf stage, with second application 10 to 14 days later if necessary. Water rate: 500 l/ha (202 l/ac).

Strawberry (Field Grown):

Non Everbearing Varieties: 3 applications of 5 l/ha (2 l/ac) from start of flowering. Repeat applications at 7 to 10 day intervals. Everbearing Varieties: Divide total rate of 15 l/ha (6 l/ac) into 3 applications of 5 l/ha (2 l/ac). Do not apply at less than 10 to 14 day intervals. Water rate: 300 to 600 l/ha (121.40-242.8 l/ac).

Sugar Beet:

3 to 5 l/ha (1.21-2 l/ac) at the 4 to 6 leaf stage. For moderate to severe deficiency, repeat at 10 to 14 day intervals. Water rate: 200 l/ha (81 l/ac).

Sweetcorn:

5 l/ha (2 l/ac) at the 4 to 6 leaf stage. Repeat at 10 to 14 day intervals if necessary. Water rate: 200 l/ha (81 l/ac).

Tobacco:

Two applications of 3 to 5 l/ha (1.21-2 l/ac) two to three weeks after transplanting (3 to 4 leaf stage) with 10 days between applications. Water rate: 400 l/ha (161.88 l/ac).

Tomatoes (Field Grown):

5 l/ha (2 l/ac) at the 4 to 6 leaf stage, with repeat applications at 10 to 14 day intervals, if necessary. Water rate: 500 l/ha (202 l/ac).

Vines:

5 l/ha (2 l/ac) at flower buds separated, fruit set, grape expansion and early veraison. Water rate: 200 l/ha (81 l/ac).